

# **Piper rhythm in the activation of the gastrocnemius medialis during running**

Lisa M. Stirling<sup>1</sup>,  
Vinzenz von Tscharnner<sup>1</sup>,  
Patrick Kugler<sup>2</sup>,  
Benno M. Nigg<sup>1</sup>

<sup>1</sup> Human Performance Laboratory, Faculty of Kinesiology, University of Calgary, Canada  
<sup>2</sup> Pattern Recognition Laboratory, University of Erlangen-Nuremberg, Germany

**For submission to the Journal of Electromyography and Kinesiology**

Corresponding Author:  
Lisa M. Stirling, BAsC, PhD  
Human Performance Laboratory, Faculty of Kinesiology  
University of Calgary  
2500 University Drive NW  
Calgary, Alberta, Canada  
T2N 1N4  
Phone: (403) 220-2170  
Fax: (403) 282-7637  
e-mail: lstirling@kin.ucalgary.ca

Running head: Piper rhythm in EMG during running

Keywords: EMG; running; time-frequency analysis; wavelet transform

## **ABSTRACT**

The presence of temporal rhythmicity in electromyographic (EMG) signals at frequencies of 35-60 Hz was initially noted by Piper (1907). This modulation and synchronization of motor unit activity is generally accepted to represent a centrally generated coding of motor commands. The purpose of this study was to resolve and quantify the Piper rhythm in the gastrocnemius medialis (GM) muscle during running. EMG was recorded from the GM of 14 female runners during 1-hour treadmill runs. The average wavelet transform was computed for EMG from series of steps taken at 2 minute intervals throughout the run. The total intensity across three wavelets (center frequencies: 170, 218, 271 Hz) was computed and a histogram indicating the incidence peaks in this signal was generated for each subject. In order to rule out effects of the analysis process, the process was repeated using simulated EMG data. Autocorrelations of the histograms were used to extract the frequency of the peaks resulting in rhythmicity at 25-55Hz. The ability to measure superimposed rhythmicity in EMG signals during dynamic tasks allows investigation of the role of aspects of central drive during movement. In particular, the changes in central control during dynamic activities can be examined with this approach.

## 1 INTRODUCTION

2 The presence of temporal rhythmicity in muscle activity has long been described during  
3 isokinetic tasks as well as during slow movements. In general, one can consider the EMG recorded  
4 from movements to comprise of muscle events (ME) and pacing events (PE). Muscle events  
5 represent a large portion of the energy in the signal and have envelopes containing low frequencies.  
6 These events correspond to the central control of the muscle providing an activation burst from  
7 onset to termination. Pacing events, on the other hand, have lower signal energy content, and  
8 represent an underlying rhythmicity contained within the muscle control signal.

9 The characteristics of the rhythms that occur during motor tasks and mechanisms proposed to  
10 explain them vary depending on the task. For example, a physiological tremor with a frequency of  
11 8-10Hz can be measured during position holding. Similarly, a discontinuity in movement occurring  
12 at a similar frequency is apparent during slow movements, although it is unclear that the  
13 mechanisms are common. The role of the spinal stretch reflex in rhythmic muscle activity has long  
14 been debated with data supporting [Hagbarth and Young, 1979; Sanes, 1985] and playing down  
15 [Jacks et al., 1988; Wessberg and Vallbo, 1995] the importance of its role. Instead the 8-10Hz  
16 rhythms were attributed to pulsatile descending signals [Wessberg and Vallbo, 1995] and the  
17 corresponding frequency modulation of active motor units [Wessberg and Kakuda, 1999].

18 In addition, motor activity has been seen to synchronise at frequencies in the beta (15-30Hz),  
19 and low gamma (30-60Hz) bands [Brown, 2000]. The specific case of modulated motor unit (MU)  
20 activity in the low gamma band is termed the Piper rhythm [Piper, 1907].

21 Rhythmicity in the Piper band has been shown to be the product of central drive, seemingly  
22 originating from the motor cortex [Brown et al., 1998]. For example, neural activity in the motor  
23 cortex has been measured and shown to be coherent with activity in various muscles [Conway et al.,  
24 1995; Salenius et al., 1996; Salenius et al., 1997; Tecchio et al., 2006]. Changes in central drive

1 relating to fatigue have been shown to manifest as a decrease in the frequency of muscle  
2 rhythmicity in the Piper band [von Tscharnner et al., 2010]. In addition, the features of muscle  
3 rhythmicity found in different pathologies may help to reveal the pathophysiology of the disorder  
4 [Brown and Marsden, 1996; McAuley et al., 2001], or conversely, clarify the pathway of the Piper  
5 rhythm [Brown, 1997]. For instance, in un-medicated Parkinson's patients, the 40-50Hz frequency  
6 associated with the Piper rhythm is replaced with a 10Hz signal indicating that the Piper pathway is  
7 influenced by activity in the pallidal projections [Brown, 1997].

8 Piper oscillations in motor output can be extracted from groups of single MU recordings  
9 [Hagbarth et al., 1983], or surface EMG signals [McAuley et al., 2001], or by measuring the  
10 resulting muscle vibrations [Brown, 1997; Hill, 1921]. In our hands, the fine structure of the  
11 rhythmic pacing events in EMG activity becomes apparent in the wavelet transformed EMG signal.  
12 Our purpose was to test the hypothesis that during running, pacing events occur with a regular  
13 timing with respect to heelstrike. In addition, the uniformity of the pacing frequency among  
14 different individuals will be investigated. A simulation of muscle activity is used to evaluate the  
15 contribution of intrinsic rhythmicity stemming from the random firing of multiple motor units and  
16 to eliminate the possibility that the observed pacing events are the result of the wavelet transform  
17 method implemented.

## 18 **1. METHODS**

19 Fifteen female recreational runners (age  $32.4 \pm 8.7$  years, mass  $61.3 \pm 6.1$  kg) participated in  
20 this study. Subjects gave their written informed consent in accordance with the university's policy  
21 on research using human subjects. The protocol was approved by the Conjoint Health Research  
22 Ethics Board at the University of Calgary.

### 23 **1.1. Measurement protocol**

1 Speed at ventilatory threshold (sVT) was determined during a preliminary session where  
2 respiratory gas exchange was monitored (Cosmed K4b2, Rome Italy) during an incremental running  
3 test on a motorized treadmill (Quinton Instrument Co., Seattle, WA, USA). Belt speed was  
4 increased by 0.13 m/s every 2 minutes until the subjects reached ventilatory threshold [Fletcher et  
5 al., 2009]. Ventilatory threshold was defined as the speed at which there was both a decrease in the  
6 fraction of expired CO<sub>2</sub> and a non-linear increase in the V<sub>E</sub>/VCO<sub>2</sub> slope [Reinhard et al., 1979;  
7 Skinner and McLellan, 1980; Wasserman et al., 1973]. Two to four weeks later, each subject  
8 returned to the lab for a 1-hr endurance running session during which they ran at approximately  
9 95% of their speed at ventilatory threshold, sVT, while electromyographic (EMG) signals were  
10 recorded from the gastrocnemius medialis (GM) muscle of the right leg.

11 The subjects were recreational athletes and were instructed to maintain their usual activity  
12 levels, therefore, significant training adaptation was not expected to occur during the interval  
13 between sessions [Gaesser and Poole, 1986]. At the selected speed of 95% of sVT the subjects were  
14 expected to be able to complete 1-hr of running while experiencing substantial levels of fatigue. In  
15 two cases 95% sVT exceeded the subjects' perceived 1-hr sustainable leg turnover (i.e., for subjects  
16 with sVT > 3.75 m/s), so the belt speed was decreased to 90% of their speed at ventilatory  
17 threshold.

18 In preparation for the run, the skin over the belly of the GM muscle was lightly abraded, cleaned  
19 with alcohol. Bipolar silver/silver chloride (Ag/AgCl) surface electrodes (Norotrode dual  
20 electrodes, Myotronics-Noromed, Inc., Kent, WA, US) were placed on the belly of the GM muscle  
21 in alignment with the direction of the muscle fibres according to SENIAM recommendations  
22 (Seniam.org). This placement ensured that the electrode remained over the surface of the muscle  
23 throughout the range of lengths corresponding to running movements. All electrodes were placed by  
24 a single experimenter to ensure consistency thorough the study. A single reference electrode was

1 secured to the tibial tuberosity. The electrodes and preamplifiers (Biovision, Wehrheim Germany)  
2 were secured to the skin using medical tape (Cover-Roll stretch, BSN medical GnbH, Hamburg,  
3 Germany) to minimize movement artefact and to prevent the electrodes from losing surface contact  
4 due to sweating. EMG signals were preamplified (1000x) and bandpass filtered (10-500 Hz) during  
5 acquisition. Heel strike was detected using an accelerometer attached externally to the heel of the  
6 right running shoe and recorded simultaneously with EMG at a sample rate of 2400 Hz. Thirty  
7 seconds of EMG and accelerometer data were recorded at 2 minute intervals throughout the run.  
8 Each 30 second data recording is referred to as a “lap”, during which approximately 40 steps  
9 occurred.

## 10 **1.2. Data Analysis**

11 All occurrences of heel strike were determined by detecting the onset of the sharp deceleration  
12 associated with floor contact. EMG representing individual steps comprised 600ms of data centered  
13 on heel strike (1440 samples). This analysis window included all potential pre- and post-heelstrike  
14 muscle activation associated with the step.

### 15 *1.2.1. Simulation of EMG activity*

16 A simulated version of the EMG was generated for every step. Simulated EMG was constructed  
17 by convolving a frequency modulated pulse train with an estimated MUAP.

18 Generation of a modulated pulse train: The 1440 sample raw EMG for each step was lowpass  
19 filtered (15Hz) to determine the rough envelop of the activity. Dividing all of the envelope’s values  
20 by the peak value resulted in a normalized envelope function. The normalized envelope was binned  
21 into 20 bins (72 samples each). The value of the bin provided a scaling factor between 0 and 1,  
22 which was used to determine the number of random pulses that were inserted into the corresponding  
23 bin in the pulse train. The maximum number of pulses per bin was set to 9 (corresponding to 300  
24 pulse/sec). If fewer than 3 pulses were required in a bin they were omitted from the pulse train. The  
25 resultant pulse train had the same length (1440 samples) as the original raw EMG signal.

1 Estimation of MUAP shape: The bin containing the maximum amplitude in the envelope function  
2 was identified for each step. The Fourier transform was used to calculate the power spectrum for the  
3 raw EMG in the corresponding bin. The power spectra from each step recorded during one lap (i.e.,  
4 30 second series of steps) were averaged. The square root of the average power spectrum was  
5 multiplied by  $i$  (square root of -1) and was submitted to the inverse Fourier transform. This yielded  
6 a symmetric waveform that mimicked the shape of the MUAP from the original data.  
7 A single estimated MUAP was generated for each lap and convolved with the pulse train generated  
8 for each individual step. This procedure resulted in unique simulated muscle activity for a given  
9 step (referred to as an “EMG step”) corresponding to each real “EMG step”.

10 This method was used to generate five sets of simulated data for each set of real data. Each set  
11 of resulting simulated data differed by the random pulse train used to generate any given EMG step.

#### 12 *1.2.2. Quantification of Pacing Events*

13 Wavelet transform: The wavelet transform of each real and simulated EMG step was performed  
14 using a set of 13 non-linearly scaled wavelets [von Tscherner, 2000]. The centre frequencies of the  
15 wavelets were: 7, 19, 38, 62, 92, 128, 170, 218, 271, 331, 395, 466 and 542 Hz. In general, the  
16 wavelet transform produces an EMG intensity pattern where time and frequency (center frequencies  
17 of the wavelet filters) are indicated on the abscissa and ordinate, respectively, and the greyscale  
18 represents the power of the transformed EMG signal.

19 The EMG average wavelet pattern was found corresponding to each lap. The average patterns  
20 from real EMG were visually inspected for movement artefact and noise content and all  
21 unacceptable average patterns were rejected along with their simulated counterparts. The total  
22 power in the band of the wavelets 7 to 9 (center frequencies 170 to 271 Hz) was found by summing  
23 the corresponding wavelet intensities for each time point. Low frequency bands were omitted from

1 the selected range due to the long time resolutions of the wavelets, and high frequency bands were  
2 omitted because of the increased amounts of power reflecting signal noise.

3 Characterization of pacing events: All peaks in the summed wavelet intensity signal were identified.  
4 Peaks occurring at less than 1% of the maximum value of the signal were neglected (i.e., small  
5 oscillations around zero). A raster plot was generated for each subject where each row indicated the  
6 indices of the peaks in the intensity data from a single lap. A histogram was generated for each  
7 subject representing the number of occurrences of peaks at each sample time (1.67ms bins).  
8 Separate raster plots and histograms were made for the real and 5 sets of simulated EMG data. Each  
9 plot was normalized to contain a total of 100 counts distributed across all bins.

10 The autocorrelation was taken of the 6 histograms generated for each subject (1 real and 5  
11 simulated EMG datasets). For each subject a net autocorrelation was calculated by subtracting the  
12 mean of the autocorrelations found from the 5 simulated datasets, from the autocorrelation found  
13 from the real data. The net autocorrelation was smoothed (low-pass 60Hz) and the timing of the first  
14 three extrema (T1, T2, T3) of the net autocorrelation were identified. The amplitude (i.e. range) of  
15 the oscillations in the net autocorrelation was determined by subtracting the mean values at the  
16 minima T1 and T3 from the value of the maximum at T2. Rhythmicity in the autocorrelation was  
17 deemed to be present if the resulting amplitude was above a threshold value.

18 The threshold value was selected to represent the minimum random oscillations expected in the  
19 autocorrelation and was determined using the autocorrelations of the 5 simulations for each  
20 individual subject. One of the 5 simulations was selected to represent the “test” data. The remaining  
21 4 simulations were used to create the average autocorrelation of the simulated data. The net  
22 autocorrelation consisted of the autocorrelation of the new “test” data minus the new average  
23 autocorrelation of the simulated data. The amplitude of the resulting net autocorrelation was found.  
24 This process was repeated 5 times for each subject, each iteration using a different simulation as the

1 “test” data. The threshold amplitude was calculated for each subject using the average of the  
2 amplitudes of the 5 resulting net autocorrelations. The corresponding threshold value was compared  
3 to each true net autocorrelation for each subject.

4 The pacing frequency for each subject with a net autocorrelation amplitude that was found to be  
5 above threshold was calculated using the equation:  $2 / (T2 + (T3-T1))$ . A linear regression was used  
6 to find the relationship between pacing frequency and amplitude. The  $R^2$  value was determined.

## 7 **2. RESULTS**

8 The average maximum aerobic speed of the 14 subjects included in this study was  $3.3 \pm 0.4$  m/s.  
9 For the test sessions the subjects ran at an average speed of 3.1 m/s ( $\pm 0.3$  m/s), representing  
10 approximately 94% ( $93.6 \pm 3.0\%$ ) of their speed at ventilatory threshold.

11 The wavelet patterns from the simulated data had a strong resemblance to those of the  
12 corresponding real data (figure 1C,D). In general, the traces of the simulated intensity bands were  
13 characterized by smaller, and more frequent oscillations (figure 1E,F). This was more apparent in  
14 the raster plot and histogram representation of the peaks in the intensity trace (figure 2).

15 **[insert figure 1]**

16 **[insert figure 2]**

17 For real EMG data, the peaks within a subject tended to have consistent timing across all laps.  
18 This resulted in a vertical alignment of dots in the raster plots, and clusters of bins with high counts  
19 in the histograms. The peaks in the simulated data were more randomly distributed. This was  
20 characterized by less structure in the raster plots, and a corresponding broad distribution of counts  
21 across all bins in the histogram. Examples of the histograms of real (top row) and simulated (middle  
22 row) data for 5 subjects are shown in figure 3, along with the corresponding autocorrelations  
23 (bottom row).

24 **[insert figure 3]**

1       Of the 14 subjects, all were found to have net autocorrelations with amplitudes above threshold.  
2       The mean ( $\pm$ std) of the threshold amplitude for significant autocorrelations was  $0.157 \pm 0.102$ . The  
3       detection of local extrema in the net autocorrelations resulted in pacing frequencies that varied from  
4       subject to subject within the range from 23.1 to 54.5 Hz (table 1). The mean ( $\pm$ std) of the pacing  
5       frequencies was  $40.6 \pm 10.7$  Hz. The relationship between pacing frequency and amplitude showed a  
6       trend towards larger amplitudes for lower frequencies. The linear regression had an  $R^2$  value of  
7       0.27.

8       **[insert table 1]**

### 9       **3. DISCUSSION**

10       There was a detectible rhythmicity within the bursts of EMG recorded from the gastrocnemius  
11       medialis during running. This pacing rhythm overlay the primary burst of muscle activity occurring  
12       after heelstrike corresponding to the generation of force required for forward propulsion during the  
13       stance phase. The frequency of the pacing rhythm was determined to range from approximately 25  
14       to 55 Hz. This band of frequencies corresponds with Piper rhythms previously measured in EMG  
15       [Brown, 2000; Piper, 1907]. It is also consistent with previous results from our group using the  
16       same method to measure rhythmicity during isometric contractions of the abductor pollicis brevis  
17       (submitted for publication).

18       Some rhythmicity was observed in the autocorrelations derived from the simulated data. In  
19       some cases this rhythmicity was consistent across all simulations for a given subject and contained  
20       peaks in the autocorrelations that were shared with those from the real data (figure 3, s9). This  
21       suggested that aspects of the general shape of the primary muscle burst were conserved in the  
22       envelop used to generate the simulated data. In other cases the rhythmicity in the simulated data was  
23       not consistent between simulations (figure 3, s1). These fluctuations in the autocorrelation represent  
24       occurrences of synchronicity arising from the random positive and negative interference of the

1 superimposed MUAPs used to generate the simulation. In both cases, subtracting the  
2 autocorrelation of the simulated data from that of the real data resulted in a net autocorrelation  
3 containing only aspects of the signal that represented the overlying pacing rhythm. This pacing  
4 rhythm was determined to be significant in all of the measured subjects.

5 Rhythms within the beta and low gamma (Piper) bands have been shown to be correlated to  
6 central descending drive [Brown, 2000]. This suggests that the rhythms that were measured from  
7 the gastrocnemius muscle during running reflect aspects of central control generated within the  
8 brain. The fact that the pacing events aligned with respect to heelstrike for all measurements taken  
9 throughout a given run (evident in rasterplots – figure 2) suggests that the heelstrike event is  
10 transmitted to higher control centers and influences the phase of the rhythmicity. Not surprisingly,  
11 the pacing varied from subject to subject both in frequency and in amplitude. This reflects varying  
12 conditions, possibly including the shoe condition, effort level or state of fatigue being experienced  
13 by the individual. Data shown in the figures of a previous study [von Tscharner et al., 2003] seemed  
14 to indicate similar pacing events occurring in the tibialis anterior. In that study, the changes seemed  
15 to be related to the footwear condition. Fatigue has specifically been shown to influence the  
16 rhythmicity at the muscle, and with respect to the brain through measurements of corticomuscular  
17 coherence [Tecchio et al., 2006; Yang et al., 2009]. The effect of these factors on the EMG pacing  
18 frequency during running will be investigated in future studies.

19 This study has some limitations, including the selection of the cut-off frequency of the lowpass  
20 filter used to extract the envelop of the real EMG signal for use in the generation of the simulated  
21 data. Despite the use of a 15Hz cut-off, it is possible that a portion of the low frequency component  
22 of the original pacing events were conserved within the envelop. However, all net autocorrelations  
23 were found to have a significant amplitude, suggesting that the selected cut-off did allow the greater  
24 portion of the pacing signal to be omitted from the simulated dataset.

## 1 **4. CONCLUSION**

2       The ability to measure superimposed rhythmicity in EMG signals during dynamic tasks allows  
3 for the investigation of the role of aspects of central drive during movement. In particular, with this  
4 method, differences in fine pacing structures overlaying the muscle events during dynamic activity  
5 can be used to investigate changes of central control in the presence external or internal influences  
6 (e.g. fatigue or footwear conditions).

7

## 1 **ACKNOWLEDGMENTS**

2 Funding was provided by the Alberta Heritage Foundation for Medical Research (AHFMR), the  
3 da Vinci Foundation and adidas AG. The sponsors had no role in the study design, in the collection,  
4 analysis and interpretation of data; in the writing of the manuscript; or in the decision to submit the  
5 manuscript for publication. The authors thank Dr. Brian MacIntosh, Jared Fletcher and Shane Esau  
6 for their contributions.

## 1 REFERENCES

- 2
- 3 Brown P, Marsden CD. Rhythmic cortical and muscle discharge in cortical myoclonus. *Brain*  
4 1996;119(4):1307-1316
- 5 Brown P. Muscle sounds in Parkinson's disease. *The Lancet* 1997;349(9051):533-535
- 6 Brown P, Salenius S, Rothwell JC, Hari R. Cortical correlate of the Piper rhythm in humans. *J*  
7 *Neurophysiol* 1998;80(6):2911-7
- 8 Brown P. Cortical drives to human muscle: the Piper and related rhythms. *Prog Neurobiol*  
9 2000;60(1):97-108
- 10 Conway BA, Halliday DM, Farmer SF, Shahani U, Maas P, Weir AI, Rosenberg JR.  
11 Synchronization between motor cortex and spinal motoneuronal pool during the performance of a  
12 maintained motor task in man. *The Journal of Physiology* 1995;489(Pt 3):917-924
- 13 Fletcher JR, Esau SP, Macintosh BR. Economy of running: beyond the measurement of oxygen  
14 uptake. *J Appl Physiol* 2009;107(6):1918-22
- 15 Gaesser GA, Poole DC. Lactate and ventilatory thresholds: disparity in time course of adaptations  
16 to training. *J Appl Physiol* 1986;61(3):999-1004
- 17 Hagbarth KE, Young RR. Participation of the stretch reflex in human physiological tremor. *Brain*  
18 1979;102(3):509-26
- 19 Hagbarth KE, Jessop J, Eklund G, Wallin EU. The Piper rhythm--a phenomenon related to muscle  
20 resonance characteristics? *Acta Physiol Scand* 1983;117(2):263-71
- 21 Hill AV. The tetanic nature of the voluntaty contraction in man. *Journal of Physiology*  
22 1921;55(sup):14-16
- 23 Jacks A, Prochazka A, Trend PS. Instability in human forearm movements studied with feed-back-  
24 controlled electrical stimulation of muscles. *J Physiol* 1988;402:443-61
- 25 McAuley JH, Corcos DM, Rothwell JC, Quinn NP, Marsden CD. Levodopa reversible loss of the  
26 Piper frequency oscillation component in Parkinson's disease. *Journal of Neurology, Neurosurgery*  
27 *& Psychiatry* 2001;70(4):471-475
- 28 Piper J. Ueber den willkuerlichen Muskeltetanus. *Pfluegers Gesamte Physiol.* 1907;119:301-338

- 1 Reinhard U, Muller PH, Schmulling RM. Determination of anaerobic threshold by the ventilation  
2 equivalent in normal individuals. *Respiration* 1979;38(1):36-42
- 3 Salenius S, Salmelin R, Neuper C, Pfurtscheller G, Hari R. Human cortical 40 Hz rhythm is closely  
4 related to EMG rhythmicity. *Neurosci Lett* 1996;213(2):75-8
- 5 Salenius S, Portin K, Kajola M, Salmelin R, Hari R. Cortical control of human motoneuron firing  
6 during isometric contraction. *J Neurophysiol* 1997;77(6):3401-5
- 7 Sanes JN. Absence of enhanced physiological tremor in patients without muscle or cutaneous  
8 afferents. *J Neurol Neurosurg Psychiatry* 1985;48(7):645-9
- 9 Skinner JS, McLellan TH. The transition from aerobic to anaerobic metabolism. *Res Q Exerc Sport*  
10 1980;51(1):234-48
- 11 Tecchio F, Porcaro C, Zappasodi F, Pesenti A, Ercolani M, Rossini PM. Cortical short-term fatigue  
12 effects assessed via rhythmic brain-muscle coherence. *Exp Brain Res* 2006;174(1):144-51
- 13 von Tscharner V. Intensity analysis in time-frequency space of surface myoelectric signals by  
14 wavelets of specified resolution. *J Electromyogr Kinesiol* 2000;10(6):433-45
- 15 von Tscharner V, Goepfert B, Nigg BM. Changes in EMG signals for the muscle tibialis anterior  
16 while running barefoot or with shoes resolved by non-linearly scaled wavelets. *Journal of*  
17 *Biomechanics* 2003;36(8):1169-76
- 18 von Tscharner V, Barandun M, Stirling LM. Fatigue-related decrease in Piper rhythm frequency of  
19 the Abductor Pollicis Brevis muscle during isometric contractions. 2010;submitted to *J*  
20 *Electromyogr Kinesiol*
- 21 Wasserman K, Whipp BJ, Koyl SN, Beaver WL. Anaerobic threshold and respiratory gas exchange  
22 during exercise. *J Appl Physiol* 1973;35(2):236-43
- 23 Wessberg J, Vallbo AB. Coding of pulsatile motor output by human muscle afferents during slow  
24 finger movements. *J Physiol* 1995;485(1):271-82
- 25 Wessberg J, Kakuda N. Single motor unit activity in relation to pulsatile motor output in human  
26 finger movements. *J Physiol* 1999;517(1):273-85
- 27 Yang Q, Fang Y, Sun C-K, Siemionow V, Ranganathan VK, Khoshknabi D, Davis MP, Walsh D,  
28 Sahgal V, Yue GH. Weakening of functional corticomuscular coupling during muscle fatigue. *Brain*  
29 *Research* 2009;1250:101-112  
30  
31  
32

## 1 **FIGURE/TABLE CAPTIONS**

2

3 Figure 1: Example of real (left) and simulated (right) EMG data from subject 4. Data is centered  
4 around heel-strike (HS) with 300ms before and after. Raw signals (A, B), wavelet intensity patterns  
5 (C, D) and the summed total intensities across the wavelet bands 7-9 (E, F) are illustrated. The  
6 white lines in the wavelet intensity patterns indicate the bottom and top of wavelet bands 7-9.

7

8 Figure 2: Example of real (left) and simulated (right) EMG data from subject 8. The raster plots (A,  
9 B) indicate the timing of peaks (x-axis) in the total intensity traces from each trial (y-axis). The  
10 histograms (C, D) indicate the number of peaks across all trials per time bin. The threshold level for  
11 peak detection is indicated by the horizontal gray line in the histograms.

12

13 Figure 3: Examples of histograms from real (top row) and simulated (middle row) EMG data from 5  
14 subjects. Resultant autocorrelations are indicated in the bottom row (upper gray – simulated data,  
15 upper black – real data, lower gray – net autocorrelation, lower black – smoothed net  
16 autocorrelation, black circles – first 2 minima (T1 and T3) and first maxima (T2) in the filtered net  
17 autocorrelation).

18

19 Table 1: Pacing frequencies (F) and amplitudes (A) extracted from the net autocorrelations of each  
20 subject. Pacing amplitude is listed in units of the autocorrelation exceeding the threshold for each  
21 individual.

## TABLES

<b>subject #</b>	<b>F [Hz]</b>	<b>A</b>
12	23.1	0.13
8	27.9	0.28
2	28.6	0.40
1	31.6	0.24
10	32.4	0.13
5	36.4	0.05
9	37.5	0.13
11	46.2	0.26
4	46.2	0.09
3	50.0	0.11
7	50.0	0.07
6	52.2	0.14
14	52.2	0.11
13	54.5	0.03

Table 1: Pacing frequencies (F) and amplitudes (A) extracted from the net autocorrelations of each subject. Subjects are ordered from lowest to highest pacing amplitude. Pacing amplitude is listed in units of the autocorrelation exceeding the threshold for each individual.

Figure 1




Figure 2





Figure 3



**Lisa M. Stirling (née Guevremont)** received her B.A.Sc. degree (with honours) in electrical engineering from the University of Toronto, Canada, in 2002. She completed her Ph.D. in medical sciences (biomedical engineering) at the University of Alberta, Canada, in 2007. Her graduate research focused on the development of control algorithms and functional electrical stimulation protocols for restoring standing and stepping after spinal cord injury. She is currently a post-doctoral fellow at the Human Performance Laboratory, University of Calgary, where she is pursuing her interests in the neural control of movement and the application of engineering approaches to the fields of rehabilitation and movement science.

**Vinzenz von Tschärner** was born in Switzerland 1947 He received his diploma in applied physics and mathematics 1974 and his PhD degree in biophysics at the University of Basel, Switzerland. He was a post doctorate fellow at Oxford University, Dep. Biochemistry, England in 1978 and 1979, and a post doctorate fellow at Stanford University, California USA Dep. Biochemistry 1998. He returned to the Biocenter in Basel 1981. He was then research affiliate at the Theodor Kocher Institute in Bern and specialized in signal transduction studying cellular responses related to cytokin binding. He became Adj. Assistant Professor (1997) and Adj. Associate Professor (2000) at the Human Performance Laboratory, University of Calgary. His main field of research is the signal propagation controlling movement patterns of humans. This involves biophysical/biomedical measurements and the analysis of sensory systems.

**Patrick F. Kugler** was born 1983 in Germany. He received his Diplom (German M.Sc.) with honours in Computer Science at the University of Erlangen-Nuremberg in 2009. During his studies he performed a half year internship at the Human Performance Lab at the University of Calgary, where he worked on his thesis on the classification of EMG signals. Currently he is working as a PhD student at the Pattern Recognition Lab at the University of Erlangen-Nuremberg. In his research he focuses on the application of algorithms from pattern recognition and machine learning to biomechanics and human motion.

**Benno M. Nigg**, was born in Switzerland, and studied nuclear physics at the world renowned ETH in Zurich, Switzerland. In 1971 switched to the study of Biomechanics. His goal was to improve individuals' mobility and longevity through first, the study of forces impacting the lower body, and then the development of orthotics, running shoes, and exercise prescriptions that would enhance the quality of individuals' lives. Dr. Nigg joined the University of Calgary as the first director of the Human Performance Laboratory in 1981. Since his arrival, he has built a team of 100 co-workers that have positioned the Human Performance Laboratory with the elite programs in the world. Dr. Nigg has published more than 280 articles in scientific journals and authored or edited ten books. He has received numerous international awards, including the prestigious Olympic Order for recognition of this outstanding service and accomplishments for the Olympic Movement.

Author Bio Photo - Stirling  
[Click here to download high resolution image](#)



Author Bio Photo - von Tscharnier  
[Click here to download high resolution image](#)



Author Bio Photo - Kugler  
[Click here to download high resolution image](#)



Author Bio Photo - Nigg  
[Click here to download high resolution image](#)

